

## TABLE OF CONTENTS

<b>CHAPTER 1 THE REGIONAL GEOLOGY AND TECTONICS OF ICELAND</b>	<b>1</b>
<b><u>Introduction</u></b>	<b>1</b>
<b><u>General Spatial and Chronological Relationships</u></b>	<b>1</b>
<b><u>Hotspots - A Review</u></b>	<b>6</b>
<b><u>Iceland Hotspot</u></b>	<b>15</b>
<b><u>Dynamics of Mantle Melting</u></b>	<b>20</b>
<b><u>Mid-Atlantic Spreading Ridge</u></b>	<b>25</b>
<b><u>Southeastern Volcanic Zone</u></b>	<b>29</b>
<b><u>South Iceland Seismic Zone</u></b>	<b>32</b>
<b><u>Mechanics of Axial Spreading</u></b>	<b>34</b>
<b>CHAPTER 2 GEOLOGY, PETROLOGY AND TECTONIC SETTING OF THE SOUTHEASTERN VOLCANIC ZONE</b>	<b>39</b>
<b><u>Introduction</u></b>	<b>39</b>
<b><u>Hekla</u></b>	<b>39</b>
<b><u>Vatnafjöll</u></b>	<b>40</b>
<b><u>Torfajökull</u></b>	<b>42</b>
<b><u>Tindfjöll</u></b>	<b>43</b>
<b><u>Katla</u></b>	<b>43</b>
<b><u>Eyjafjöll Volcanic System</u></b>	<b>44</b>
<b><u>Vestmannæyar</u></b>	<b>46</b>
<b><u>Chemical, Petrologic, and Geophysical Data for the SEVZ</u></b>	<b>47</b>
<b>Isotopic Compositions</b>	<b>47</b>
<b>Petrology</b>	<b>51</b>
<b>Geochemistry</b>	<b>51</b>
<b>Geophysical data</b>	<b>53</b>
<b><u>Geochemical Models and Trends for the Propagating Rift</u></b>	<b>53</b>
<b><u>Research Objectives</u></b>	<b>64</b>

<b><u>CHAPTER 3</u></b>	<b>METHODS OF ANALYSIS</b>	<b>66</b>
	<b><u>K-Ar Radiometric Dating</u></b>	<b>66</b>
	<b><u>Microprobe Analysis</u></b>	<b>67</b>
	<b><u>XRF Analysis</u></b>	<b>68</b>
	<b><u>INAA Analysis</u></b>	<b>69</b>
	<b><u>Helium Isotopic Analyses</u></b>	<b>72</b>
	<b><u>Modeling Programs</u></b>	<b>74</b>
<b><u>CHAPTER 4</u></b>	<b>PETROGRAPHIC, GEOCHEMICAL AND GEOCHRONOLOGICAL DATA FOR EYJAFJÖLL</b>	<b>76</b>
	<b><u>Stratigraphy</u></b>	<b>76</b>
	<b><u>Major and Trace Element Data</u></b>	<b>81</b>
	<b><u>Petrography</u></b>	<b>95</b>
	<b>Groundmass</b>	<b>95</b>
	<b>Vesicles</b>	<b>98</b>
	<b>Phenocrysts</b>	<b>98</b>
	<b>PLAGIOCLASE</b>	<b>99</b>
	<b>CLINOPYROXENE</b>	<b>99</b>
	<b>OLIVINE</b>	<b>102</b>
	<b>TITANOMAGNETITE</b>	<b>102</b>
	<b>CUMULATE ROCKS</b>	<b>105</b>
	<b><u>Mineral Compositions</u></b>	<b>106</b>
	<b>PLAGIOCLASE</b>	<b>106</b>
	<b>CLINOPYROXENE</b>	<b>110</b>
	<b>OLIVINE</b>	<b>110</b>
	<b>TITANOMAGNETITE</b>	<b>110</b>
	<b>GLASS</b>	<b>113</b>
	<b><u>Isotopic Compositions</u></b>	<b>113</b>
<b><u>CHAPTER 5</u></b>	<b>INTERPRETATION AND DISCUSSION OF GEOCHEMISTRY, GEOCHRONOLOGY, AND TECTONICS OF THE EYJAFJÖLL VOLCANIC SYSTEM</b>	<b>114</b>
	<b><u>Geochemistry</u></b>	<b>114</b>
	<b>Mineral Data - Magma Dynamics</b>	<b>114</b>
	<b>Chemical Data - Crystallization Models</b>	<b>119</b>

Oxygen Fugacity, Crystallization Sequence, and Pressure	131
Magma Chambers	134
REE Data - Discussion	137
Helium Data - Crustal Assimilation?	144
Formation of Alkalic Magmas	147
<u>Geochronology</u>	152
Stratigraphy	152
Chemical Evolution of Eyjafjöll Volcanic System Through Time	153
<u>Tectonic Implications</u>	159
Trends Along the SEVZ	159
Propagating Rifts and the SEVZ	163
Tectonic Models for the SEVZ	169
 <u>CHAPTER 6 CONCLUSIONS AND FUTURE WORK</u>	 171
<u>Conclusions</u>	171
<u>Future Work</u>	173
 <b>BIBLIOGRAPHY</b>	 176
 APPENDICES	
<u>Appendix A</u> HANDSAMPLE AND LOCATION DESCRIPTIONS	184
<u>Appendix B</u> GEOCHEMICAL DATA	192
<u>Appendix C</u> THIN SECTION PETROGRAPHY	207
<u>Appendix D</u> MICROPROBE DATA	216

## LIST OF FIGURES

<b>Figure 1.01:</b> Map of Iceland, showing Eyjafjöll (topped by a glacier), Vestmannæyar, and Reykjavik.	2
<b>Figure 1.02:</b> Map of Iceland showing the distribution of volcanism according to age, from the Tertiary to present.	3
<b>Figure 1.03:</b> Map of Iceland showing the active volcanic and tectonic zones: Reykjanes Ridge (RR), Kolbeinsey Ridge (KR), Tjörnes Fracture Zone (TFZ), Northern Rift Zone (NRZ), Eastern Rift Zone (ERZ), Southeastern Volcanic Zone (SEVZ), Western Rift Zone (WRZ), Snæfellsnes Volcanic Zone (SVZ), Mid-Iceland Transform Zone (MITZ), South Iceland Seismic Zone (SISZ).	5
<b>Figure 1.04:</b> Block diagram illustrating the topographic expression of various types of dilational structures seen in the active volcanic zones of Iceland.	7
<b>Figure 1.05:</b> A photograph of a laboratory starting convective plume showing the structure caused by conduction of heat and consequent entrainment of surrounding fluid.	9
<b>Figure 1.06:</b> A model of the evolution of Sr ratios with time, assuming that half the Earth's mantle has had its low-melting constituents extracted to give the equivalent of a 50 km thick crust, (from Hall, 1987).	11
<b>Figure 1.07:</b> Plot of Sr ratios against Nd ratios for basaltic lava suites, (from Hall, 1987).	12
<b>Figure 1.08:</b> Graphical representation of the Holmes-Houtermans model.	14
<b>Figure 1.09:</b> Isotope ratios of Pb from young volcanic rocks.	14
<b>Figure 1.10:</b> Reconstruction of the North Atlantic at magnetic anomaly C23, just after the onset of sea-floor spreading.	18
<b>Figure 1.11:</b> The morphology of the Greenland-Scotland ridge (from Bott, 1981).	19

<b>Figure 1.12:</b> Cross section lines for figures 1.13 and 1.14.	21
<b>Figure 1.13:</b> Cross-sections spanning latitude range from 63.0°N to 66.0°N, from NW to SE (A-A') and from SW to NE (B-B').	22
<b>Figure 1.14:</b> Generalized crust and upper mantle cross-section derived from the RRISP main profile.	22
<b>Figure 1.15:</b> Major and trace element variation within Iceland (Elliott et al., 1991).	24
<b>Figure 1.16:</b> Pb and Nd variation within Iceland.	24
<b>Figure 1.17:</b> Schematic representation of normal ridge (A) and propagating ridge (B) transform intersections.	26
<b>Figure 1.18:</b> Southeastern Volcanic Zone (SEVZ), with the volcanic systems (dotted fields) of Hekla, Vatnafjöll, Torfajökull, Tindfjöll, Eyjafjöll, Katla, and Vestmannæyar (after Jakobsson, 1979).	30
<b>Figure 1.19:</b> The South Iceland Seismic Zone (SISZ) and the Southeastern Volcanic Zone (SEVZ) showing mapped surface breaks and regions in which over half of the buildings were destroyed in historic seismic events.	31
<b>Figure 1.20:</b> Bookshelf faulting.	33
<b>Figure 1.21:</b> Depth to the lower crust (layer 3) in Iceland (in km).	35
<b>Figure 1.22:</b> Development of the present geometry of the tectonically active zones in Iceland.	36
<b>Figure 1.23:</b> Some major identified magnetic anomaly lineations in the ocean surrounding Iceland (from Björnsson, 1981).	37
<b>Figure 2.01:</b> The silica contents of the initial gas-rich products of historic eruptions from Hekla tend to be higher than those of the main basaltic magma that follows, and the amount of this difference is directly proportional to the time of repose between eruptions.	41

<b>Figure 2.02:</b> Eyjafjöll Volcanic System - Geology (modified from Jakobsson, 1979, and Jón Jónsson, 1988 ).	45
<b>Figure 2.03:</b> $^3\text{He}/^4\text{He}$ values for different regions within Iceland.	48
<b>Figure 2.04:</b> La/Sm histograms for different volcanic regions.	49
<b>Figure 2.05:</b> $^{87}\text{Sr}/^{86}\text{Sr}$ histograms for different regions within Iceland.	49
<b>Figure 2.06:</b> Pb-Pb diagrams for Recent and Tertiary volcanics from Iceland.	50
<b>Figure 2.07:</b> Nd ratios versus Sr ratios for both Recent and Tertiary volcanics from Iceland (from Meyer et al., 1985).	50
<b>Figure 2.08:</b> Mineralogical variations along the southern neovolcanic zones.	52
<b>Figure 2.09:</b> The location of drillholes used to determine thermal gradients in Iceland.	54
<b>Figure 2.10:</b> Geothermal gradients in Iceland and their relationship to the pyrolite solidus.	55
<b>Figure 2.11:</b> P-T diagram for partial melting of pyrolite, indicating compositions of liquids generated.	60
<b>Figure 2.12:</b> Schematic diagram of southern volcanic zones in Iceland illustrating magma genesis and evolution (from Meyer et al., 1985).	61
<b>Figure 2.13:</b> Model of magma generation and accumulation along the SEVZ.	63
<b>Figure 4.01:</b> Eyjafjöll Volcanic System.	77
<b>Figure 4.02:</b> Eyjafjöll Volcanic System - Geology (modified from Jakobsson, 1979, and Jón Jónsson, 1988 ).	78

<b>Figure 4.03:</b> Approximate stratigraphy of lowest portion of Steinafjall sampling section.	82
<b>Figure 4.04:</b> Geological cross-section of Hvammsmúli from the East, from Steinthórsson (1964).	83
<b>Figure 4.05 (a-m):</b> Major (a-h) and trace (i-m) element variation diagrams, with data for the Eyjafjöll Volcanic System.	84
<b>Figure 4.06:</b> Eyjafjöll data according to the IUGS classification from LeBas et al. (1986).	91
<b>Figure 4.07:</b> Alkali vs. silica diagram for Eyjafjöll samples.	92
<b>Figure 4.08:</b> AFM diagram for Eyjafjöll data with tholeiite and alkaline trends from Irvine and Baragar (1972).	93
<b>Figure 4.09:</b> A segment of the basalt tetrahedron with Eyjafjöll data (all samples with SiO <sub>2</sub> < 55 wt%).	94
<b>Figure 4.10:</b> FeO vs. TiO <sub>2</sub> for all Eyjafjöll samples with SiO <sub>2</sub> < 55 wt%.	96
<b>Figure 4.11:</b> Chondrite-normalized REE patterns for Eyjafjöll samples.	97
<b>Figure 4.12:</b> Thin section of Ey-10 showing plagioclase phenocryst with different zoning geometries (XPL).	100
<b>Figure 4.13:</b> Thin section of Ey-29 showing plagioclase phenocryst with small clinopyroxene and opaque inclusions.	101
<b>Figure 4.14:</b> Thin section of Ey-39 showing two clinopyroxene phenocrysts growing together (PPL).	103
<b>Figure 4.15:</b> Thin section of Ey-36 showing olivine phenocryst with remnants of rounded glass inclusion (0.4 mm across).	104
<b>Figure 4.16:</b> Microprobe analysis from core to rim of a plagioclase phenocryst from Ey-10.	109

<b>Figure 4.17:</b>	111
Pyroxene triangle showing averages of clinopyroxene phenocryst compositions for all Eyjafjöll samples.	
<b>Figure 4.18:</b>	112
Thin section of Ey-1 showing titanomagnetite phenocryst (reflected light).	
<b>Figure 5.01:</b>	116
Representative olivine phenocrysts from Ey-17, showing three different core compositions.	
<b>Figure 5.02:</b>	118
Representative plagioclase phenocrysts from Ey-10, showing multiple core compositions and zoning types.	
<b>Figure 5.03 (a-m):</b>	120
Major (a-h) and trace (i-m) element variation diagrams, including data for the Eyjafjöll Volcanic System, and trends for a perfect fractional crystallization model (MIXNFRAC) at the NNO buffer and QFM buffer.	
<b>Figure 5.04:</b>	130
Change in density during crystallization of parent magma (Ey-49) at the NNO buffer.	
<b>Figure 5.05:</b>	135
Ti vs. Al (cation mole fraction) for experimental clinopyroxenes from Vestmannæyar study (Thy, 1991a).	
<b>Figure 5.06:</b>	136
Ti vs. Al (cation mole fraction) for clinopyroxenes from Eyjafjöll data (full range of Ti data).	
<b>Figure 5.07:</b>	136
Ti vs. Al (cation mole fraction) for clinopyroxenes from Eyjafjöll data (limited range of Ti data).	
<b>Figure 5.08:</b>	138
Chondrite-normalized REE patterns for Eyjafjöll samples.	
<b>Figure 5.09:</b>	141
Chondrite-normalized REE patterns for Ey-4 and STF-3 (representing the highest and lowest abundances for Eyjafjöll).	
<b>Figure 5.10:</b>	142
Nb and Zr variations for Eyjafjöll data combined with modeled fractional crystallization trend.	
<b>Figure 5.11:</b>	143
Incompatible element patterns for Eyjafjöll samples, normalized to normal MORB (NMORB) concentrations.	



<b>Figure 5.12:</b> Compositions for mantle melts at different starting compositions and pressures (depths of melting).	149
<b>Figure 5.13:</b> Model of magma generation and accumulation along the SEVZ.	151
<b>Figure 5.14:</b> A segment of the basalt tetrahedron with Eyjafjöll data sorted according to age.	154
<b>Figure 5.15:</b> Alkali vs. silica diagram with Eyjafjöll data sorted according to age.	155
<b>Figure 5.16:</b> Accumulation rates for lavas on Eyjafjöll throughout known volcanic history.	158
<b>Figure 5.17:</b> Alkali vs. silica diagram for data from Eyjafjöll, with fields corresponding to data from Hekla, Katla, and Vestmannæyar.	160
<b>Figure 5.18:</b> A segment of the basalt tetrahedron with data from Eyjafjöll and fields corresponding to data from Hekla, Katla, and Vestmannæyar.	161
<b>Figure 5.19:</b> Chondrite-normalized REE patterns for samples from Eyjafjöll, the Hekla-Katla region, and Vestmannæyar.	164
<b>Figure 5.20:</b> Idealized cooling paths for crustal magma bodies.	165

## LIST OF TABLES

<b>Table 1.01:</b> Buoyancy, volume, and magma volume fluxes for specific hotspot localities throughout the world.	17
<b>Table 3.01:</b> Precision of XRF analysis.	70
<b>Table 3.02:</b> Accuracy of XRF analysis.	71
<b>Table 3.03:</b> Standard analyses and associated error for INAA accuracy.	73
<b>Table 4.01:</b> Eyjafjöll stratigraphy.	79
<b>Table 4.02:</b> Microprobe data.	107
<b>Table 4.03:</b> Representative analyses from microprobe analysis.	108
<b>Table 5.01:</b> Chemical effects of magma mixing and crystal accumulation.	127
<b>Table 5.02:</b> Compositions of a spinel lherzolite (Kushiro et al., 1968), most evolved possible parent magma from MIXNFRAC fractional crystallization model (liquid just before plagioclase crystallization begins, 1187°C), and most primitive possible parent magma, from same model (Ey-49).	150

## LIST OF APPENDICES FIGURES

<b>Figure D.01 (a-f):</b> Microprobe transects of plagioclase phenocrysts.	219
<b>Figure D.02 (a-g):</b> Microprobe transects of olivine phenocrysts.	226

## LIST OF APPENDICES TABLES

<b>Table B.01:</b> Major and trace element data from XRF analysis.	193
<b>Table B.02:</b> CIPW norms, and equilibrium olivine data.	200
<b>Table B.03:</b> Ta, Hf, and REE data from INAA.	205
<b>Table B.04:</b> K-Ar ages and $^3\text{He}/^4\text{He}$ values.	206
<b>Table C.01:</b> Thin section petrography.	208
<b>Table D.01:</b> Representative analyses of plagioclase phenocrysts from Microprobe Analysis.	217
<b>Table D.02:</b> Representative analyses of olivine phenocrysts from Microprobe Analysis.	223

## LIST OF PLATES

### PLATE 1:

MACINTOSH DISK WITH MICROPROBE DATA  
(Data stored in Microsoft Excel 2.2 spreadsheets.)